


Anti-Aquaporin 2 Antibody (A89116)

Specifications:

| Description:Rabbit polyclonal antibody to Aquaporin 2.Applications:WB, IHC, ICC/IFRecommended Dilutions:WB: 1:500-1:1,000, IHC: 1:50-1:100, ICC/IF: 1:50-1:200Reactivity:Human, Mouse, Rat |
|--|
| Recommended Dilutions:WB: 1:500-1:1,000, IHC: 1:50-1:100, ICC/IF: 1:50-1:200Reactivity:Human, Mouse, Rat |
| Reactivity: Human, Mouse, Rat |
| |
| |
| Immunogen:A synthetic peptide corresponding to a sequence within amino acids 100-200 of humanAQP2 (NP_000477.1). |
| Sequence: GAALLHEITPADIRGDLAVNALSNSTTAGQAVTVELFLTLQLVLCIFASTDERRGENP GTPALSIGFSVALGHLLGIHYTGCSMNPARSLAPAVVTGKFDD |
| Host: Rabbit |
| Clonality: Polyclonal |
| Isotype: IgG |
| Conjugate: Unconjugated |
| Purification: Affinity purification. |
| Molecular Weight: 26 kDa |
| Product Form: Liquid |
| Formulation: Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.05% Proclin 300. |
| Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use. |

Images:




Western blot analysis of extracts of 293T cells, using Anti-Aquaporin 2 Antibody (A89116) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 90s.


antibodies

Anti-Aquaporin 2 Antibody (A89116)


Images continued:



Immunohistochemistry analysis of paraffin-embedded mouse kidney using Anti-Aquaporin 2 Antibody (A89116) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunofluorescence analysis of rat kidney cells using Anti-Aquaporin 2 Antibody (A89116) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of mouse kidney cells using Anti-Aquaporin 2 Antibody (A89116) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).