


## Anti-ILF3 Antibody (A9053)

### Specifications:


Name:	Anti-ILF3 Antibody
Description:	Rabbit polyclonal antibody to ILF3.
Applications:	WB, IHC, ICC/IF, IP
Recommended Dilutions:	WB: 1:500-1:2,000, IHC: 1:50-1:200, ICC/IF: 1:50-1:200, IP: 1:500-1:1,000
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-270 of human ILF3 (NP_036350.2).
Sequence:	MRP_MRIFVNDDR_HVMAKHSSVYPTQEELEAVQNMVSHTERALKAVSDWIDEQEKGSS EAESDNMDVPPEDDSKEGAGEQKTEHMTRTLRGV_MRVGLVAKGLLLKGDLDLELVLLC KEKPTTALLDKVADNLAIQLAAVTEDKYEILQSVDDAAIVIKNTKEPPLSLTIHLTSP VVREEMEKVLAGETLSVNDPPDVLDRQKCLAALASLRHAKWFQARANGLKSCVIVIRV LRDLCTR_VPTWGPLRGWPLELLCEKSIGTANRPMGAGE
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	95 / 120 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

## Anti-ILF3 Antibody (A9053)


### Images:




Western blot analysis of extracts of various cell lines, using Anti-ILF3 Antibody (A9053) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 $\mu$ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 1s.



Western blot analysis of extracts of mouse thymus, using Anti-ILF3 Antibody (A9053) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 $\mu$ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 5s.



Immunofluorescence analysis of C6 cells using Anti-ILF3 Antibody (A9053) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of NIH-3T3 cells using Anti-ILF3 Antibody (A9053) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).