


Anti-Nanog Antibody (A9165)

Specifications:

Name:	Anti-Nanog Antibody
Description:	Rabbit polyclonal antibody to Nanog.
Applications:	WB, IHC
Recommended Dilutions:	WB: 1:1,000-1:2,000, IHC: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-105 of human Nanog (NP_079141.2).
Sequence:	MSVDPACPQSLPCFEASDCKESSPMPVICGPEENYPQLQSSAEMPHTETVSPPLPSSM DLLIQDSPDSSTSPKGGKQPTSAEKSVAKKEDKVPVKKQKTRTVFSST
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	48 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.


Images:



Western blot analysis of extracts of NIH/3T3 cells, using Anti-Nanog Antibody (A9165) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 45s.

Anti-Nanog Antibody (A9165)

Images continued:



Immunohistochemistry analysis of paraffin-embedded human testicular seminoma using Anti-Nanog Antibody (A9165) at a dilution of 1:20 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.