


Anti-MMP9 Antibody (A91306)

Specifications:


Name:	Anti-MMP9 Antibody
Description:	Rabbit polyclonal antibody to MMP9.
Applications:	IHC, ICC/IF
Recommended Dilutions:	IHC: 1:50-1:200, ICC/IF: 1:50-1:200
Reactivity:	Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 20-707 of human MMP9 (NP_004985.2).
Sequence:	APRQRQSTLVLPFGDLRTNLTDRQLAEEYLYRYGYTRVAEMRGESKSLGPALLLLQKQ LSLPETGELDSATLKAMRTPRCGVPDLGRFQTFEGDLKWHHHNITYWIQNYSEDLPRA VIDDAFARAFALWSAVTPLTFTRVYSRDADIVIQFGVAEHGDGYFPDGDGLLAHAFP PGPGIQGDAHFDDELWSLGKGVVVPTRFGNADGAACHFPFIFEGRSYSACTTDGRSD GLPWCSTTANYDTDDRFGFCPSERLYTQDGNADGKPCQFPFIFQGGQSYSACTTDGRSD GYRWCATTANYDRDKLFGFCPTRADSTVMGGNSAGELCVFPFTFLGKEYSTCTSEGRG DGRLWCATTSNFDSDDKKGWFCPDQGYSLFLVAAHEFGHALGLDHSSVPEALMYPMYRF TEGPPLHKDDVNGIRHLYGPRPEPEPRPPTTTTPQPTAPPTVCPTGPPTVHPSERPTA GPTGPPSAGPTGPPTAGPSTATTVPLSPVDDACNVNIFDAIAEIGNQLYLFKDGKYWR FSEGRGSRPQGPFLIADKWPALPRKLDVFEERLSKLLFFFSGRQVWVYTGASVLGPR RLDKLGLGADVAQVTGALRSGRGKMLLFSGRRLWRFDVKAQMVDPRSASEVDRMFPGV PLDTHDVFQYREKAYFCQDRFYWRVSSRSELNQVDQVGYVTYDILQCPED
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	92 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-MMP9 Antibody (A91306)


Images:




Immunohistochemistry analysis of paraffin-embedded rat spleen using Anti-MMP9 Antibody (A91306) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse testis using Anti-MMP9 Antibody (A91306) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse spleen using Anti-MMP9 Antibody (A91306) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunofluorescence analysis of L929 cells using Anti-MMP9 Antibody (A91306) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).