


Anti-MUC2 Antibody (A91628)

Specifications:


Name:	Anti-MUC2 Antibody
Description:	Rabbit polyclonal antibody to MUC2.
Applications:	WB, IHC, ICC/IF
Recommended Dilutions:	WB: 1:500-1:1,000, IHC: 1:100-1:500, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 21-200 of human MUC2 (NP_002448.4).
Sequence:	SELQTEGRTRNHGHNV CSTWGNFHYKTFDGDVFRFPGLCDYNFASDCRGSYKEFAVHL KRGPGQAEAPAGVESILLTIKDDTIYLTRHLAVLNGAVVSTPHYSPGLLIEKSDAYTK VYSRAGLTLMWNREDALMLELDTKFRNHTCGLCGDYNGLQSYSEFLSDGVLFSPLEFG NMQKIN
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	110 kDa / 130 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.05% Proclin 300.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-MUC2 Antibody (A91628)


Images:




Western blot analysis of extracts of HT-29 cells, using Anti-MUC2 Antibody (A91628) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Enhanced Kit (RM00021). Exposure time: 180s.



Western blot analysis of extracts of various cell lines, using Anti-MUC2 Antibody (A91628) at 1:500 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Enhanced Kit (RM00021). Exposure time: 180s.



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue using Anti-MUC2 Antibody (A91628) at a dilution of 1:300 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human colon tissue using Anti-MUC2 Antibody (A91628) at a dilution of 1:300 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.