

Anti-FADD Antibody (A92876)


Specifications:

Name:	Anti-FADD Antibody
Description:	Rabbit polyclonal antibody to FADD.
Applications:	WB
Recommended Dilutions:	WB: 1:500-1:2,000
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-208 of human FADD (Q13158).
Sequence:	MDPFLVLLHSVSSSLSSSELTELKFLCLGRVGKRKLERVQSGLDLFSMLLEQNDLEPG HTELLRELLASLRRHDLLRRVDDFEAGAAAGAAPGEEDLCAAFNVICDNVGKDWRRLA RQLKVSDTKIDSIEDRYPRNLTERVRESLRIWKNTEKENATVAHLVGALRSCQMNLVA DLVQEVQQARDLQNRSGAMSPMSWNSDASTSEAS
Host:	Rabbit
Clonality:	Polyclonal
Clonality: Isotype:	Polyclonal IgG
-	•
Isotype:	lgG
lsotype: Conjugate:	IgG Unconjugated
Isotype: Conjugate: Purification:	IgG Unconjugated Affinity purification.
Isotype: Conjugate: Purification: Molecular Weight:	IgG Unconjugated Affinity purification. 28 kDa
Isotype: Conjugate: Purification: Molecular Weight: Product Form:	IgG Unconjugated Affinity purification. 28 kDa Liquid
Isotype: Conjugate: Purification: Molecular Weight: Product Form: Formulation:	IgG Unconjugated Affinity purification. 28 kDa Liquid Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.



Anti-FADD Antibody (A92876)

Images:



Western blot analysis of extracts from normal (control) and FADD knockout (KO) HeLa cells, using Anti-FADD Antibody (A92876) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 5s.