

Anti-FADD Antibody (A14994)


Specifications:

Name:	Anti-FADD Antibody
Description:	Rabbit polyclonal antibody to FADD.
Applications:	WB, IP
Recommended Dilutions:	WB: 1:500-1:2,000, IP: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-208 of human FADD (NP_003815.1).
Sequence:	MDPFLVLLHSVSSSLSSSELTELKFLCLGRVGKRKLERVQSGLDLFSMLLEQNDLEPG HTELLRELLASLRRHDLLRRVDDFEAGAAAGAAPGEEDLCAAFNVICDNVGKDWRRLA RQLKVSDTKIDSIEDRYPRNLTERVRESLRIWKNTEKENATVAHLVGALRSCQMNLVA DLVQEVQQARDLQNRSGAMSPMSWNSDASTSEAS
Host:	Rabbit
Host: Clonality:	Rabbit Polyclonal
Clonality:	Polyclonal
Clonality: Isotype:	Polyclonal IgG
Clonality: Isotype: Conjugate:	Polyclonal IgG Unconjugated
Clonality: Isotype: Conjugate: Purification:	Polyclonal IgG Unconjugated Affinity purification.
Clonality: Isotype: Conjugate: Purification: Molecular Weight:	Polyclonal IgG Unconjugated Affinity purification. 26 kDa
Clonality: Isotype: Conjugate: Purification: Molecular Weight: Product Form:	Polyclonal IgG Unconjugated Affinity purification. 26 kDa Liquid
Clonality: Isotype: Conjugate:	Polyclonal IgG Unconjugated
Clonality: Isotype: Conjugate: Purification: Molecular Weight:	Polyclonal IgG Unconjugated Affinity purification. 26 kDa
Clonality: Isotype: Conjugate: Purification: Molecular Weight: Product Form:	Polyclonal IgG Unconjugated Affinity purification. 26 kDa Liquid
Clonality: Isotype: Conjugate: Purification: Molecular Weight: Product Form: Formulation:	Polyclonal IgG Unconjugated Affinity purification. 26 kDa Liquid Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.


antibodies

Anti-FADD Antibody (A14994)

Images:



Western blot analysis of extracts of various cell lines, using Anti-FADD Antibody (A14994) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit.



Immunoprecipitation analysis of 200µg extracts of HeLa cells using 3µg of Anti-FADD Antibody (A14994). This Western blot was performed on the immunoprecipitate using Anti-FADD Antibody (A14994) at a dilution of 1:1000.