


## Anti-TLR6 Antibody (A10330)

## Specifications:

| Name:                  | Anti-TLR6 Antibody   |
|------------------------|--|
| Description:           | Rabbit polyclonal antibody to TLR6.  |
| Applications:          | WB, ICC/IF   |
| Recommended Dilutions: | WB: 1:500-1:1,000, ICC/IF: 1:50-1:200  |
| Reactivity:            | Human, Mouse   |
| Immunogen:             | A synthetic peptide corresponding to a sequence within amino acids 697-796 of human TLR6 (NP_006059.2).  |
| Sequence:              | KSIFVLSPNFVQSEWCHYELYFAHHNLFHEGSNNLILILLEPIPQNSIPNKYHKLKAL<br>MTQRTYLQWPKEKSKRGLFWANIRAAFNMKLTLVTENNDVKS |
| Host:                  | Rabbit   |
| Clonality:             | Polyclonal   |
| lsotype:               | lgG  |
| Conjugate:             | Unconjugated   |
| Purification:          | Affinity purification.   |
| Molecular Weight:      | 100 kDa  |
| Product Form:          | Liquid   |
| Formulation:           | Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.05% Proclin 300.                  |
| Storage:               | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.                    |
| Disclaimer:            | This product is for research use only. It is not intended for diagnostic or therapeutic use.             |
|                        |  |

## Images:




Western blot analysis of extracts of various cell lines, using Anti-TLR6 Antibody (A10330) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 10s.



## Anti-TLR6 Antibody (A10330)

Images continued:



Immunofluorescence analysis of Jurkat cells using Anti-TLR6 Antibody (A10330) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).