


Anti-Noxa Antibody (A10802)

Specifications:

| | |
|------------------------|--|
| Name: | Anti-Noxa Antibody |
| Description: | Rabbit polyclonal antibody to Noxa. |
| Applications: | WB, ICC/IF |
| Recommended Dilutions: | WB: 1:500-1:2,000, ICC/IF: 1:50-1:200 |
| Reactivity: | Human, Mouse, Rat |
| Immunogen: | A synthetic peptide corresponding to a sequence within amino acids 1-54 of human PMAIP1 (NP_066950.1). |
| Sequence: | MPGKKARKNAQPSPARAPAELEVECATQLRRFGDKLNFRQKLLNLISKLFCSGT |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Isotype: | IgG |
| Conjugate: | Unconjugated |
| Purification: | Affinity purification. |
| Molecular Weight: | 14 kDa |
| Product Form: | Liquid |
| Formulation: | Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium Azide. |
| Storage: | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| Disclaimer: | This product is for research use only. It is not intended for diagnostic or therapeutic use. |


Images:




Western blot analysis of extracts of various cell lines, using Anti-Noxa Antibody (A10802) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 90s.

Anti-Noxa Antibody (A10802)


Images continued:



Immunofluorescence analysis of Jurkat cells using Anti-Noxa Antibody (A10802) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of rat thymus cells using Anti-Noxa Antibody (A10802) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of mouse thymus cells using Anti-Noxa Antibody (A10802) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).