


Anti-S6K1 (phospho Thr389) Antibody (A11042)

Specifications:

Name:	Anti-S6K1 (phospho Thr389) Antibody
Description:	Rabbit polyclonal antibody to S6K1 (phospho Thr389).
Applications:	WB, ICC/IF
Recommended Dilutions:	WB: 1:100-1:500, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	A synthetic phosphorylated peptide around T389 of human P70 S6K (NP_003152.1).
Sequence:	GFTYV
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	70 kDa / 85 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.05% Proclin 300.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.


Images:




Western blot analysis of NIH/3T3, using Anti-S6K1 (phospho Thr389) Antibody (A11042) at 1:1,000 dilution. NIH/3T3 cells were treated by 10% FBS at 37°C for 30 minutes after serum-starvation overnight. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 180s.

Anti-S6K1 (phospho Thr389) Antibody (A11042)

Images continued:



Immunofluorescence analysis of C6 cells using Anti-S6K1 (phospho Thr389) Antibody (A11042) at a dilution of 1:100. C6 cells were treated by Serum-starvation overnight at 37°C.



Immunofluorescence analysis of HeLa cells using Anti-S6K1 (phospho Thr389) Antibody (A11042) at a dilution of 1:100. HeLa cells were treated by Serum-starvation overnight at 37°C.