

## Anti-DUSP14 Antibody (A11277)


### Specifications:

| Name:                  | Anti-DUSP14 Antibody   |
|------------------------|--|
| Description:           | Rabbit polyclonal antibody to DUSP14.  |
| Applications:          | WB, ICC/IF   |
| Recommended Dilutions: | WB: 1:200-1:2,000, ICC/IF: 1:50-1:200  |
| Reactivity:            | Human, Mouse, Rat  |
| Immunogen:             | Recombinant fusion protein containing a sequence corresponding to amino acids 1-198 of human DUSP14 (NP_008957.1).   |
| Sequence:              | MSSRGHSTLPRTLMAPRMISEGDIGGIAQITSSLFLGRGSVASNRHLLQARGITCIVN<br>ATIEIPNFNWPQFEYVKVPLADMPHAPIGLYFDTVADKIHSVSRKHGATLVHCAAGVS<br>RSATLCIAYLMKFHNVCLLEAYNWVKARRPVIRPNVGFWRQLIDYERQLFGKSTVKMV<br>QTPYGIVPDVYEKESRHLMPYWGI |
| Host:                  | Rabbit   |
| Clonality:             | Polyclonal   |
| Isotype:               | lgG  |
| Conjugate:             | Unconjugated   |
| Purification:          | Affinity purification.   |
| Molecular Weight:      | 22 kDa   |
| Product Form:          | Liquid   |
| Formulation:           | Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium Azide.   |
| Storage:               | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.  |
| Disclaimer:            | This product is for research use only. It is not intended for diagnostic or therapeutic use.   |


# antibodies

### Anti-DUSP14 Antibody (A11277)

#### Images:



Western blot analysis of extracts of various cell lines, using Anti-DUSP14 Antibody (A11277) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Enhanced Kit (RM00021). Exposure time: 30s.



Immunofluorescence analysis of H9C2 cells using Anti-DUSP14 Antibody (A11277) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).