

Anti-CDKN2AIP Antibody (A14412)


Specifications:

| Name: | Anti-CDKN2AIP Antibody |
|------------------------|---|
| Description: | Rabbit polyclonal antibody to CDKN2AIP. |
| Applications: | WB, IHC |
| Recommended Dilutions: | WB: 1:500-1:1,000, IHC: 1:50-1:100 |
| Reactivity: | Human, Mouse, Rat |
| Immunogen: | Recombinant fusion protein containing a sequence corresponding to amino acids 1-220 of human CDKN2AIP (NP_060102.1). |
| Sequence: | MAQEVSEYLSQNPRVAAWVEALRCDGETDKHWRHRRDFLLRNAGDLAPAGGAASASTD EAADAESGTRNRQLQQLISFSMAWANHVFLGCRYPQKVMDKILSMAEGIKVTDAPTYT TRDELVAKVKKRGISSSNEGVEEPSKKRVIEGKNSSAVEQDHAKTSAKTERASAQQEN SSTCIGSAIKSESGNSARSSGISSQNSSTSDGDRSVSSQSSSVSS |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| lsotype: | lgG |
| Conjugate: | Unconjugated |
| Purification: | Affinity purification. |
| Molecular Weight: | 65 kDa |
| Product Form: | Liquid |
| Formulation: | Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium Azide. |
| Storage: | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| Disclaimer: | This product is for research use only. It is not intended for diagnostic or therapeutic use. |
| | |

antibodies

Anti-CDKN2AIP Antibody (A14412)

Images:



Western blot analysis of extracts of various cell lines, using Anti-CDKN2AIP Antibody (A14412) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 5s.