

Anti-APPBP1 Antibody (A14542)


Specifications:

Name:	Anti-APPBP1 Antibody
Description:	Rabbit polyclonal antibody to APPBP1.
Applications:	WB
Recommended Dilutions:	WB: 1:500-1:1,000
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-220 of human NAE1 (NP_003896.1).
Sequence:	MAQLGKLLKEQKYDRQLRLWGDHGQEALESAHVCLINATATGTEILKNLVLPGIGSFT IIDGNQVSGEDAGNNFFLQRSSIGKNRAEAAMEFLQELNSDVSGSFVEESPENLLDND PSFFCRFTVVVATQLPESTSLRLADVLWNSQIPLLICRTYGLVGYMRIIIKEHPVIES HPDNALEDLRLDKPFPELREHFQSYDLDHMEKKDHSHTPWIVIIAK
Llast	B.443
Host:	Rabbit
Clonality:	Polyclonal
Clonality:	Polyclonal
Clonality: Isotype:	Polyclonal IgG
Clonality: Isotype: Conjugate:	Polyclonal IgG Unconjugated
Clonality: Isotype: Conjugate: Purification:	Polyclonal IgG Unconjugated Affinity purification.
Clonality: Isotype: Conjugate: Purification: Molecular Weight:	Polyclonal IgG Unconjugated Affinity purification. 60 kDa
Clonality: Isotype: Conjugate: Purification: Molecular Weight: Product Form:	Polyclonal IgG Unconjugated Affinity purification. 60 kDa Liquid
Clonality: Isotype: Conjugate: Purification: Molecular Weight: Product Form: Formulation:	Polyclonal IgG Unconjugated Affinity purification. 60 kDa Liquid Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.

antibodies

Anti-APPBP1 Antibody (A14542)

Images:



Western blot analysis of extracts of various cell lines, using Anti-APPBP1 Antibody (A14542) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 30s.