

## **Anti-ADAM15 Antibody (A15874)**

## Specifications:

Name: Anti-ADAM15 Antibody

Description: Rabbit polyclonal antibody to ADAM15.

Applications: WB, ICC/IF

Recommended Dilutions: WB: 1:500-1:2,000, ICC/IF: 1:50-1:200

Reactivity: Human, Mouse, Rat

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 540-700

of human ADAM15 (NP\_003806.3).

Sequence: QPAAPLCLQTANTRGNAFGSCGRNPSGSYVSCTPRDAICGQLQCQTGRTQPLLGSIRD

LLWETIDVNGTELNCSWVHLDLGSDVAQPLLTLPGTACGPGLVCIDHRCQRVDLLGAQ

ECRSKCHGHGVCDSNRHCYCEEGWAPPDCTTQLKATSSLTTGLLL

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Conjugate: Unconjugated

Purification: Affinity purification.

Molecular Weight: 93 kDa

Product Form: Liquid

Formulation: Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium

Azide.


Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.




## Anti-ADAM15 Antibody (A15874)

## Images:



Western blot analysis of extracts of various cell lines, using Anti-ADAM15 Antibody (A15874) at 1:500 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 $\mu$ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 90s.



Immunofluorescence analysis of HeLa cells using Anti-ADAM15 Antibody (A15874) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).