Anti-YAP1 Antibody (A16747)

Specifications:

|	
Name:	Anti-YAP1 Antibody
Description:	Rabbit polyclonal antibody to YAP1.
Applications:	WB, IHC, ICC/IF, IP
Recommended Dilutions:	WB: 1:500-1:2,000, IHC: 1:50-1:200, ICC/IF: 1:50-1:200, IP: 1:500-1:1,000
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 155-504 of human YAP1 (NP_001123617.1).
Sequence:	PTAQHLRQSSFEIPDDVPLPAGWEMAKTSSGQRYFLNHIDQTTTWQDPRKAMLSQMNV TAPTSPPVQQNMMNSASGPLPDGWEQAMTQDGEIYYINHKNKTTSWLDPRLDPRFAMN QRISQSAPVKQPPPLAPQSPQGGVMGGSNSNQQQQMRLQQLQMEKERLRLKQQELLRQ AMRNINPSTANSPKCQELALRSQLPTLEQDGGTQNPVSSPGMSQELRTMTTNSSDPFL NSGTYHSRDESTDSGLSMSSYSVPRTPDDFLNSVDEMDTGDTINQSTLPSQQNRFPDY LEAIPGTNVDLGTLEGDGMNIEGEELMPSLQEALSSDILNDMESVLAATKLDKESFLT WL
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	72 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-YAP1 Antibody (A16747)

Images:

Immunofluorescence analysis of C6 cells using Anti-YAP1 Antibody (A16747) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).

Immunofluorescence analysis of L929 cells using Anti-YAP1 Antibody (A16747) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).

Confocal immunofluorescence analysis of U-2 OS cells using Anti-YAP1 Antibody (A16747) at a dilution of 1:200. DAPI was used to stain the cell nuclei (blue).

Immunoprecipitation analysis of 300µg extracts of HeLa cells using 3µg of Anti-YAP1 Antibody (A16747). This Western blot was performed on the immunoprecipitate using Anti-YAP1 Antibody (A16747) at a dilution of 1:1000.