


Goat Anti-Mouse IgG H&L Antibody (Cyanine 3) (A17349)


Specifications:

| Name: | Goat Anti-Mouse IgG H&L Antibody (Cyanine 3) |
|------------------------|---|
| Description: | Goat Anti-Mouse IgG H&L Antibody (Cyanine 3). |
| Host Species: | Goat |
| Target Species: | Mouse |
| Applications: | ICC/IF, Flow Cytometry |
| Recommended Dilutions: | Flow Cytometry: 1:100-1:800, ICC/IF: 1:50-1:200 |
| Immunogen: | Mouse IgG. |
| Clonality: | Polyclonal |
| lsotype: | lgG |
| Conjugate: | Cyanine 3 |
| Product Form: | Liquid |
| Purification: | Affinity purification. |
| Formulation: | Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.75% BSA, and 0.025% Sodium Azide. |
| Storage: | Store in the dark at 4°C. Avoid prolonged exposure to light. Do not freeze! |
| Disclaimer: | This product is for research use only. It is not intended for diagnostic or therapeutic use. |

Images:



Immunofluorescence analysis of NIH-3T3 cells using Goat Anti-mouse IgG H&L Antibody (Cyanine 3) (A17349) at a dilution of 1:100/1:200 (40x lens). DAPI was used to stain the cell nuclei (blue).




Immunofluorescence analysis of PC-12 cells using Goat Anti-mouse IgG H&L Antibody (Cyanine 3) (A17349) at a dilution of 1:100/1:200 (40x lens). DAPI was used to stain the cell nuclei (blue).



Goat Anti-Mouse IgG H&L Antibody (Cyanine 3) (A17349)

Images continued:



Immunofluorescence analysis of HeLa cells using Goat Anti-mouse IgG H&L Antibody (Cyanine 3) (A17349) at a dilution of 1:100/1:200 (40x lens). DAPI was used to stain the cell nuclei (blue).