

Rabbit Anti-Goat IgG H&L Antibody (Cyanine 3) (A17354)

Specifications:

Rabbit Anti-Goat IgG H&L Antibody (Cyanine 3) Name:

Description: Rabbit Anti-Goat IgG H&L Antibody (Cyanine 3).

Host Species: Rabbit

Target Species: Goat

Applications: ICC/IF, Flow Cytometry

ICC/IF: 1:100-1:800, Flow Cytometry: 1:100-1:800 Recommended Dilutions:

Immunogen: Goat IgG.

Polyclonal Clonality:

Isotype: **IgG**

Conjugate: Cyanine 3

Product Form: Liquid

Affinity purification. Purification:


Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.75% BSA, and Formulation:

0.025% Sodium Azide.

Storage: Store in the dark at 4°C. Avoid prolonged exposure to light. Do not freeze!

Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.

Images:




Western blot analysis of extracts of 293T cells, using Rabbit Anti-Goat IgG H&L Antibody (Cyanine 3) (A17354) at 1:2,000 dilution. 293T cells were treated by 10% FBS at 37°C for 30 minutes after serum-starvation overnight. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25μg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL

Basic Kit. Exposure time: 10s.



Rabbit Anti-Goat IgG H&L Antibody (Cyanine 3) (A17354)

Images continued:



Immunohistochemistry analysis of paraffin-embedded human kidney using Rabbit Anti-Goat IgG H&L Antibody (Cyanine 3) (A17354) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.