


Anti-p63 Antibody [TP63/1786] (A250429)

Specifications:


| | |
|------------------------|--|
| Name: | Anti-p63 Antibody [TP63/1786] |
| Description: | Mouse monoclonal [TP63/1786] antibody to p63. |
| Applications: | ELISA, WB |
| Recommended Dilutions: | WB: 1-2 µg/ml |
| Reactivity: | Human |
| Immunogen: | Recombinant fragment, around amino acids 3-106, of human p63 protein. The exact sequence is proprietary. |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Clone ID: | TP63/1786 |
| Isotype: | IgG2b |
| Light Chains: | kappa |
| Conjugate: | Unconjugated |
| Purification: | Protein A/G chromatography. |
| Concentration: | 200 µg/ml |
| Product Form: | Liquid |
| Formulation: | Supplied in 10mM Phosphate Buffered Saline with 0.05% BSA and 0.05% Sodium Azide. |
| Storage: | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| General Notes: | This monoclonal antibody is also available in a different formulation without BSA and Sodium Azide - Anti-p63 Antibody [TP63/1786] - BSA and Azide free (A253609). |
| Disclaimer: | This product is for research use only. It is not intended for diagnostic or therapeutic use. |

Anti-p63 Antibody [TP63/1786] (A250429)


Images:



Western blot analysis of recombinant p63 protein, PC3 cell lysate, and HeLa cell lysate using Anti-p63 Antibody [TP63/1786].




Western blot analysis of MCF-7, T47D, and MDA-MB453 cell lysates using Anti-p63 Antibody [TP63/1786].



SDS-PAGE analysis of Anti-p63 Antibody [TP63/1786] under non-reduced and reduced conditions; showing intact IgG and intact heavy and light chains, respectively. SDS-PAGE analysis confirms the integrity and purity of the antibody.

Anti-p63 Antibody [TP63/1786] (A250429)

Images continued:



Analysis of protein array containing more than 19,000 full-length human proteins using Anti-p63 Antibody [TP63/1786]. Z-Score and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target; a MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.