


Anti-Ferritin Light Chain Antibody [FTL/1386] - BSA and Azide free (A251828)

Specifications:


| | |
|------------------------|--|
| Name: | Anti-Ferritin Light Chain Antibody [FTL/1386] - BSA and Azide free |
| Description: | Mouse monoclonal [FTL/1386] antibody to Ferritin Light Chain. |
| Applications: | ELISA, WB, IHC |
| Recommended Dilutions: | WB: 1-2 µg/ml, IHC: 0.25-0.5 µg/ml |
| Reactivity: | Human |
| Immunogen: | Recombinant fragment, around amino acids 38-165, of human FTL protein. The exact sequence is proprietary. |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Clone ID: | FTL/1386 |
| Isotype: | IgG2b |
| Light Chains: | kappa |
| Conjugate: | Unconjugated |
| Purification: | Protein A/G chromatography. |
| Concentration: | 1 mg/ml |
| Product Form: | Liquid |
| Formulation: | Supplied in 10mM Phosphate Buffered Saline; without Sodium Azide and carrier free. |
| Storage: | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| General Notes: | This monoclonal antibody is also available in a different formulation with BSA and Sodium Azide - Anti-Ferritin Light Chain Antibody [FTL/1386] (A248646). |
| Disclaimer: | This product is for research use only. It is not intended for diagnostic or therapeutic use. |

Anti-Ferritin Light Chain Antibody [FTL/1386] - BSA and Azide free (A251828)


Images:



Western blot analysis of A431, HeLa, liver, and testis lysates using Anti-Ferritin Light Chain Antibody [FTL/1386].




Western blot analysis of human HeLa cell lysate using Anti-Ferritin Light Chain Antibody [FTL/1386].



SDS-PAGE analysis of Anti-Ferritin Light Chain Antibody [FTL/1386] under non-reduced and reduced conditions; showing intact IgG and intact heavy and light chains, respectively. SDS-PAGE analysis confirms the integrity and purity of the antibody.

Anti-Ferritin Light Chain Antibody [FTL/1386] - BSA and Azide free (A251828)

Images continued:



Analysis of protein array containing more than 19,000 full-length human proteins using Anti-Ferritin Light Chain Antibody [FTL/1386]. Z-Score and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target; a MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.