


Anti-c-Kit Antibody [KIT/2673] - BSA and Azide free (A252290)

Specifications:


| | |
|------------------------|---|
| Name: | Anti-c-Kit Antibody [KIT/2673] - BSA and Azide free |
| Description: | Mouse monoclonal [KIT/2673] antibody to c-Kit. |
| Specificity: | This antibody recognizes a protein of 145kDa, identified as CD117/p145kit. It is found on a wide variety of tumor cells including follicular and papillary carcinoma of thyroid, adenocarcinomas from endometrium, lung, ovary, pancreas, and breast as well as malignant melanoma, endodermal sinus tumor, and small cell carcinoma. However, anti-CD117 has been particularly useful in differentiating gastrointestinal stromal tumors from Kaposis sarcoma, tumors of smooth muscle origin, fibromatosis, and neural tumors of the GI tract. Anti-CD117 is also useful in recognizing myeloblasts in bone marrow biopsy and clot section. |
| Applications: | ELISA, Flow Cytometry, WB |
| Recommended Dilutions: | Flow Cytometry: 1-2 µg/million cells, WB: 1-2 µg/ml |
| Reactivity: | Human |
| Immunogen: | Recombinant full-length human KIT protein. |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Clone ID: | KIT/2673 |
| Isotype: | IgG1 |
| Light Chains: | kappa |
| Conjugate: | Unconjugated |
| Purification: | Protein A/G chromatography. |
| Concentration: | 1 mg/ml |
| Product Form: | Liquid |
| Formulation: | Supplied in 10mM Phosphate Buffered Saline; without Sodium Azide and carrier free. |
| Storage: | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| General Notes: | This monoclonal antibody is also available in a different formulation with BSA and Sodium Azide - Anti-c-Kit Antibody [KIT/2673] (A249110). |
| Disclaimer: | This product is for research use only. It is not intended for diagnostic or therapeutic use. |

Anti-c-Kit Antibody [KIT/2673] - BSA and Azide free (A252290)

Images:



SDS-PAGE analysis of Anti-c-Kit Antibody [KIT/2673] under non-reduced and reduced conditions; showing intact IgG and intact heavy and light chains, respectively. SDS-PAGE analysis confirms the integrity and purity of the antibody.



Analysis of protein array containing more than 19,000 full-length human proteins using Anti-c-Kit Antibody [KIT/2673]. Z-Score and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target; a MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.