

## Anti-TDP2 Antibody [TDP2/1258] - BSA and Azide free (A252771)

### Specifications:

Name: Anti-TDP2 Antibody [TDP2/1258] - BSA and Azide free

Description: Mouse monoclonal [TDP2/1258] antibody to TDP2.

Specificity: This antibody recognizes a protein of 41kDa, which is identified as TDP2. It is a member of

a superfamily of divalent cation-dependent phosphodiesterases. The encoded protein associates with CD40, tumor necrosis factor (TNF) recentor-75 and TNF recentor

associates with CD40, tumor necrosis factor (TNF) receptor-75 and TNF receptor associated factors (TRAFs), and inhibits nuclear factor-kappa-B activation. This protein has

sequence and structural similarities with APE1 endonuclease, which is involved in both DNA repair and the activation of transcription factors. DNA repair enzyme that can remove a variety of covalent adducts from DNA through hydrolysis of a 5'-phosphodiester bond,

giving rise to DNA with a free 5' phosphate. Catalyzes the hydrolysis of dead-end complexes between DNA and the topoisomerase 2 (TOP2) active site tyrosine residue.

Hydrolyzes 5'-phosphoglycolates on protruding 5' ends on DNA double-strand breaks

(DSBs) due to DNA damage by radiation and free radicals. The 5'-tyrosyl DNA

phosphodiesterase activity can enable the repair of TOP2-induced DSBs without the need for nuclease activity, creating a 'clean' DSB with 5'-phosphate termini that are ready for ligation. Has also 3'-tyrosyl DNA phosphodiesterase activity, but less efficiently and much slower than TDP1. May also act as a negative regulator of ETS1 and may inhibit nuclear

factor-kappa-B activation.

Applications: Flow Cytometry, IF, IHC-P

Recommended Dilutions: Flow Cytometry: 1-2 μg/million cells, IF: 1-2 μg/ml, IHC-P: 1-2 μg/ml

Reactivity: Human

Immunogen: Recombinant full-length human TDP2 protein.

Host: Mouse

Clonality: Monoclonal

Clone ID: TDP2/1258

Isotype: IgG2b

Light Chains: kappa

Conjugate: Unconjugated

Purification: Protein A/G chromatography.

Concentration: 1 mg/ml

Product Form: Liquid

Formulation: Supplied in 10mM Phosphate Buffered Saline; without Sodium Azide and carrier free.



# Anti-TDP2 Antibody [TDP2/1258] - BSA and Azide free (A252771)

### Specifications continued:


Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

General Notes: This monoclonal antibody is also available in a different formulation with BSA and Sodium


Azide - Anti-TDP2 Antibody [TDP2/1258] (A249591).

Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.


### Images:




Immunohistochemical analysis of formalin-fixed, paraffin-embedded human lung carcinoma using Anti-TDP2 Antibody [TDP2/1258].



Immunohistochemical analysis of formalin-fixed, paraffin-embedded human prostate carcinoma using Anti-TDP2 Antibody [TDP2/1258].



Immunohistochemical analysis of formalin-fixed, paraffin-embedded human duodenal carcinoma using Anti-TDP2 Antibody [TDP2/1258].



Immunohistochemical analysis of formalin-fixed, paraffin-embedded human colon carcinoma using Anti-TDP2 Antibody [TDP2/1258].