

Anti-CD27 Antibody [LPFS2/1611] - BSA and Azide free (A253799)

Specifications:

Name: Anti-CD27 Antibody [LPFS2/1611] - BSA and Azide free

Description: Mouse monoclonal [LPFS2/1611] antibody to CD27.

Specificity: This antibody recognizes a protein of a disulfide-linked 120kDa dimer, identified as CD27. It

is expressed on the majority of peripheral T cells, medullary thymocytes, memory-type B cells, and natural killer cells. It is a transmembrane phosphoglycoprotein that belongs to the tumor necrosis factor receptor (TNFR) superfamily. CD27 binds to its ligand CD70, a member of the TNF family, and induces T-cell co-stimulation and B-cell activation. It also interacts with TRAFs and is involved in activation of NFB and SAPK/JNK and induces

apoptosis.

Applications: ELISA, Flow Cytometry, IF, IHC-P

Recommended Dilutions: Flow Cytometry: 1-2 µg/million cells, IF: 1-2 µg/ml, IHC-P: 1-2 µg/ml

Reactivity: Human

Immunogen: Recombinant full-length human CD27 protein.

Host: Mouse

Clonality: Monoclonal

Clone ID: LPFS2/1611

Isotype: IgG1

Light Chains: kappa

Conjugate: Unconjugated

Purification: Protein A/G chromatography.

Concentration: 1 mg/ml

Product Form: Liquid

Formulation: Supplied in 10mM Phosphate Buffered Saline; without Sodium Azide and carrier free.

Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

General Notes: This monoclonal antibody is also available in a different formulation with BSA and Sodium


Azide - Anti-CD27 Antibody [LPFS2/1611] (A250619).

Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.




Anti-CD27 Antibody [LPFS2/1611] - BSA and Azide free (A253799)


Images:




Immunohistochemical analysis of formalin-fixed, paraffin-embedded human tonsil using Anti-CD27 Antibody [LPFS2/1611].



SDS-PAGE analysis of Anti-CD27 Antibody [LPFS2/1611] under non-reduced and reduced conditions; showing intact IgG and intact heavy and light chains, respectively. SDS-PAGE analysis confirms the integrity and purity of the antibody.



Immunohistochemical analysis of formalin-fixed, paraffin-embedded human tonsil using Anti-CD27 Antibody [LPFS2/1611].



Immunohistochemical analysis of formalin-fixed, paraffin-embedded human spleen using Anti-CD27 Antibody [LPFS2/1611].