


## Anti-SIRT3 Antibody [PCRP-SIRT3-1C10] (A277615)

### Specifications:


Name:	Anti-SIRT3 Antibody [PCRP-SIRT3-1C10]
Description:	Mouse monoclonal [PCRP-SIRT3-1C10] antibody to SIRT3.
Specificity:	Zinc-fingers and homeobox (ZHX) proteins are transcription factors that interact with the activation domain of the A subunit of nuclear factor-Y (NF-YA). ZHX1-3 are ubiquitously expressed proteins expressed in various tissues. They act as transcriptional repressors and localize to the nucleus. The ZHX proteins contain two Cys2-His2-type zinc-finger motifs and five homeodomains (HDs). These domains allow the ZHX proteins to form homodimers, but they can also form heterodimers with each other. However, this dimerization is not required for repressor activity. Hypermethylation-mediated silencing of ZHX2 is an epigenetic event involved in hepatocellular carcinoma (HCC).
Applications:	ELISA, IP, Flow Cytometry, IF, WB
Recommended Dilutions:	IP: 1-2µg / 100-500µg proteins, Flow Cytometry: 1-2 µg/million cells, IF: 1-2 µg/ml, WB: 1-2 µg/ml
Reactivity:	Human
Immunogen:	Recombinant full-length human SIRT3 protein.
Host:	Mouse
Clonality:	Monoclonal
Clone ID:	PCRP-SIRT3-1C10
Isotype:	IgG2a
Conjugate:	Unconjugated
Purification:	Protein A/G chromatography.
Concentration:	200 µg/ml
Product Form:	Liquid
Formulation:	Supplied in 10mM Phosphate Buffered Saline with 0.05% BSA and 0.05% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
General Notes:	This monoclonal antibody is also available in a different formulation without BSA and Sodium Azide - Anti-SIRT3 Antibody [PCRP-SIRT3-1C10] - BSA and Azide free (A278203).
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

## Anti-SIRT3 Antibody [PCRP-SIRT3-1C10] (A277615)


### Images:



Flow cytometric analysis of PFA-fixed HeLa cells using Anti-SIRT3 Antibody [PCRP-SIRT3-1C10] followed by Goat Anti-Mouse IgG (CF® 488) (Blue). Isotype Control (Red).




SDS-PAGE analysis of Anti-SIRT3 Antibody [PCRP-SIRT3-1C10] under non-reduced and reduced conditions; showing intact IgG and intact heavy and light chains, respectively. SDS-PAGE analysis confirms the integrity and purity of the antibody.



Immunofluorescent analysis of PFA-fixed HeLa cells stained with Anti-SIRT3 Antibody [PCRP-SIRT3-1C10] followed by Goat Anti-Mouse IgG (CF® 488) (Blue). Isotype Control (Red).

## Anti-SIRT3 Antibody [PCRP-SIRT3-1C10] (A277615)

Images continued:



Analysis of protein array containing more than 19,000 full-length human proteins using Anti-SIRT3 Antibody [PCRP-SIRT3-1C10]. Z-Score and S-Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target; a MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.