

Anti-alpha 1 Antichymotrypsin Antibody [SERPINA3/4189] - BSA and Azide free (A278146)

0		- : c:		4: -	
	ne	CITI	ca	TIC	ns
$\overline{}$	\sim	· · · ·	-		,,,,

Name: Anti-alpha 1 Antichymotrypsin Antibody [SERPINA3/4189] - BSA and Azide free

Description: Mouse monoclonal [SERPINA3/4189] antibody to alpha 1 Antichymotrypsin.

Specificity: It recognizes a protein of 65-76kDa, which is identified antichymotrypsin (AACT). AACT is a

plasma protease inhibitor synthesized in the liver as a single glycopeptide chain. In human, the normal serum level of AACT is about one-tenth that of α 1-antitrypsin (AAT), with which

it shares nucleic acid and protein sequence homology. Both are major acute phase reactants; their concentrations in plasma increase in response to trauma, surgery and infection. Elevated levels of AACT are widely, but not universally, reported in the cerebrospinal fluid and plasma of AD patients. Prostate-specific antigen (PSA) and its SDS-stable complex with AACT are in widespread use as markers for the diagnosis of prostate cancer. AACT deficiency may also be a possible cause of chronic liver disease.

AACT antibody reacts with histiocytes and histiocytic neoplasms. It is widely used to identify histiocytes and tumors derived from them. Acinar tumors of the pancreas and salivary gland

may also exhibit AACT positivity.

Applications: IHC-P

Recommended Dilutions: IHC-P: 1-2 μg/ml

Reactivity: Human

Immunogen: Recombinant fragment, around amino acids 49-187, of human alpha 1 Antichymotrypsin.

The exact sequence is proprietary.

Host: Mouse

Clonality: Monoclonal

Clone ID: SERPINA3/4189

Isotype: IgG2a

Light Chains: kappa

Conjugate: Unconjugated

Purification: Protein A/G chromatography.

Concentration: 1 mg/ml

Product Form: Liquid

Formulation: Supplied in 10mM Phosphate Buffered Saline; without Sodium Azide and carrier free.



Anti-alpha 1 Antichymotrypsin Antibody [SERPINA3/4189] - BSA and Azide free (A278146)

Specifications continued:


Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

General Notes: This monoclonal antibody is also available in a different formulation with BSA and Sodium


Azide - Anti-alpha 1 Antichymotrypsin Antibody [SERPINA3/4189] (A277558).

Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.

Images:



Immunohistochemical analysis of formalin-fixed, paraffin-embedded human liver tissue using Anti-alpha 1 Antichymotrypsin Antibody [SERPINA3/4189].



SDS-PAGE analysis of Anti-alpha 1 Antichymotrypsin Antibody [SERPINA3/4189] under non-reduced and reduced conditions; showing intact IgG and intact heavy and light chains, respectively. SDS-PAGE analysis confirms the integrity and purity of the antibody.



Anti-alpha 1 Antichymotrypsin Antibody [SERPINA3/4189] - BSA and Azide free (A278146)

Images continued:



Analysis of protein array containing more than 19,000 full-length human proteins using Anti-alpha 1 Antichymotrypsin Antibody [SERPINA3/4189]. Z-Score and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target; a MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.