

Anti-HSP27 Antibody (A305135)

Specifications:

Name: Anti-HSP27 Antibody

Description: Rabbit polyclonal antibody to HSP27.

Applications: WB, ICC/IF, Flow Cytometry, FACS, IHC

Recommended Dilutions: WB: 1:2,000, ICC/IF: 1:250

Reactivity: Human, Mouse, Shark, Canine

Immunogen: Human HSP27, His tagged.

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Conjugate: Unconjugated

Purification: Protein A purification.

Concentration: 1.38 mg/ml

Molecular Weight: ~27 kDa

Product Form: Liquid


Formulation: Supplied in Phosphate Buffered Saline, pH 7.4, with 50% Glycerol and 0.09% Sodium

Azide.

Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.

Images:



Immunohistochemistry analysis of human skeletal muscle, fixed in formalin and paraffin-embedded. The Primary Antibody used was Anti-HSP27 Antibody (A305135).




Anti-HSP27 Antibody (A305135)

Images continued:



Immunocytochemistry/Immunofluorescence analysis of human heat shocked cervical cancer cell line (HeLa), fixed in 2% formaldehyde for 20 minutes at room temperature, using Anti-HSP27 Antibody (A305135), at 1:250 for 12 hours at 4°C. The secondary antibody used was APC Goat Anti-Rabbit (red) at 1:200 for 2 hours at room temperature. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at room temperature. Localization: Cytoplasm. Mitochondrion matrix. Magnification: 20x.(A) DAPI (blue) nuclear stain. (B) Anti-Hsp27 Antibody. (C) Composite. Heat Shocked at 42°C for 1h.



Western blot analysis of human Cervical cancer cell line (HeLa) lysate showing detection of HSP27 protein using Anti-HSP27 Antibody (A305135) at 1:2000 for 2 hours at room temperature. Load: 15 µgprotein. Block: 1.5% BSA. The secondary antibody used was Donkey Anti-Rabbit IgG: HRP for 1 hour at room temperature.



Immunocytochemistry/Immunofluorescence analysis of human heat shocked cervical cancer cell line (HeLa), fixed in 2% formaldehyde for 20 minutes at room temperature, using Anti-HSP27 Antibody (A305135), at 1:250 for 12 hours at 4°C. The secondary antibody used was FITC Goat Anti-Rabbit (green) at 1:200 for 2 hours at room temperature. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at room temperature. Localization: Cytoplasm. Mitochondrion matrix. Magnification: 100x.(A) DAPI (blue) nuclear stain. (B) Anti-Hsp27 Antibody. (C) Composite. Heat Shocked at 42°C for 1h.