


Anti-Axl Antibody [ARC53006] (A305703)

Specifications:


| | |
|------------------------|--|
| Name: | Anti-Axl Antibody [ARC53006] |
| Description: | Rabbit monoclonal [ARC53006] antibody to Axl. |
| Applications: | WB |
| Recommended Dilutions: | WB: 1:1,000-1:5,000 |
| Reactivity: | Human, Mouse |
| Immunogen: | Recombinant fusion protein containing a sequence corresponding to amino acids 33-449 of human Axl (NP_068713.2). |
| Sequence: | EESPFVGNPGNITGARGLTGTLRCQLQVQGEPPVHWRDGGQILELADSTQTQVPLGE DEQDDWIVVSQLRITSLQLSDTGQYQCLVFLGHQTFVSQPGYVGLGLPYFLEEPEDR TVAANTPFNLSCQAQGPPEPVDLLWLQDAVPLATAPGHGPPQRSLHVPGLNKTSSFSC AHNAKGVTTSTRATITVLPQQPRNLHLVSRQPTELEVAVWTPGLSGIYPLTHCTLQAVL SDDGMGIQAGEPDPPEEPLTSQASVPPHQLRLGSLHPHTPYHIRVACTSSQGPSSWTH WLPVETPEGVPLGPPENISATRNGSQAFVHWQEPRAPLQGTLLGYRLAYQQGQDTPEVL MDIGLRQEVTLELQGDGVSNSLTVCAAYTAAGDGPWSLPVPLEAWRPGQAQPVHQLV KEPSTPAFSWP |
| Host: | Rabbit |
| Clonality: | Monoclonal |
| Clone ID: | ARC53006 |
| Isotype: | IgG |
| Conjugate: | Unconjugated |
| Purification: | Affinity purification. |
| Molecular Weight: | 138 kDa |
| Product Form: | Liquid |
| Formulation: | Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.05% Proclin 300. |
| Storage: | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| Disclaimer: | This product is for research use only. It is not intended for diagnostic or therapeutic use. |

Anti-Axl Antibody [ARC53006] (A305703)

Images:



Western blot analysis of HeLa, using Anti-Axl Antibody [ARC53006] (A305703) at 1:3,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 30s.



Western blot analysis of Mouse heart, using Anti-Axl Antibody [ARC53006] (A305703) at 1:3,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 180s.