

## **Anti-USP39 Antibody [ARC2588] (A306442)**

## Specifications:

Name: Anti-USP39 Antibody [ARC2588]

Description: Rabbit monoclonal [ARC2588] antibody to USP39.

Applications: WB, IHC, ICC/IF

Recommended Dilutions: WB: 1:500-1:1,000, IHC: 1:50-1:200, ICC/IF: 1:50-1:200

Reactivity: Human, Mouse, Rat

Immunogen: A synthetic peptide corresponding to a sequence within amino acids 466-565 of human

USP39 (Q53GS9).

Sequence: KNPTIVNFPITNVDLREYLSEEVQAVHKNTTYDLIANIVHDGKPSEGSYRIHVLHHGT

GKWYELQDLQVTDILPQMITLSEAYIQIWKRRDNDETNQQGA

Host: Rabbit

Clonality: Monoclonal

Clone ID: ARC2588

Isotype: IgG

Conjugate: Unconjugated

Purification: Affinity purification.

Molecular Weight: 65 kDa

Product Form: Liquid

Formulation: Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02%

Sodium Azide.


Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.




## **Anti-USP39 Antibody [ARC2588] (A306442)**


## Images:




Western blot analysis of extracts of Mouse liver, using Anti-USP39 Antibody [ARC2588] (A306442) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 1s.



Immunohistochemistry analysis of paraffin-embedded human lung cancer using Anti-USP39 Antibody [ARC2588] (A306442) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse lung using Anti-USP39 Antibody [ARC2588] (A306442) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse spleen using Anti-USP39 Antibody [ARC2588] (A306442) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.