


Anti-XIAP Antibody [ARC51953] (A307914)

Specifications:


Name:	Anti-XIAP Antibody [ARC51953]
Description:	Rabbit monoclonal [ARC51953] antibody to XIAP.
Applications:	WB, IP
Recommended Dilutions:	WB: 1:2,000-1:6,000, IP: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-200 of human XIAP (NP_001158.2).
Sequence:	MTFNSFEGSKTCVPADINKEEEFVEEFNRLKTFANFPSGSPVSASTLARAGFLYTGEG DTRVRCFSCHAAVDRWQYGDSAVGRHRKRVSPNCRFINGFYLENSATQSTNSGIQNGQYK VENYLGSRDHFALDRPSETHADYLLRTGQVVDISDIYPRNPAMYSEEARLKSFQNW DYAHLTPRELASAGLYYTGIGDQVQC
Host:	Rabbit
Clonality:	Monoclonal
Clone ID:	ARC51953
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	53 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.05% Proclin 300.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-XIAP Antibody [ARC51953] (A307914)

Images:



Western blot analysis of various lysates, using Anti-XIAP Antibody [ARC51953] (A307914) at 1:5,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 180s.



Western blot analysis of extracts from normal (control) and XIAP knockout (KO) HeLa(KO) cells, using Anti-XIAP Antibody [ARC51953] (A307914) at 1:5,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 180s.