

Anti-HDAC6 Antibody [ARC0805] (A307941)

Specifications:

Name: Anti-HDAC6 Antibody [ARC0805]

Description: Rabbit monoclonal [ARC0805] antibody to HDAC6.

Applications: WB, IHC, IP

Recommended Dilutions: WB: 1:500-1:1,000, IHC: 1:50-1:200, IP: 1:500-1:1,000

Reactivity: Human

Immunogen: A synthetic peptide corresponding to a sequence within amino acids 1-100 of human

HDAC6 (Q9UBN7).

Sequence: MTSTGQDSTTTRQRRSRQNPQSPPQDSSVTSKRNIKKGAVPRSIPNLAEVKKKGKMKK

LGQAMEEDLIVGLQGMDLNLEAEALAGTGLVLDEQLNEFHCL

Host: Rabbit

Clonality: Monoclonal

Clone ID: ARC0805

Isotype: IgG

Conjugate: Unconjugated

Purification: Affinity purification.

Molecular Weight: 160 kDa

Product Form: Liquid

Formulation: Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02%

Sodium Azide.


Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.




Anti-HDAC6 Antibody [ARC0805] (A307941)


Images:



Western blot analysis of extracts of various cell lines, using Anti-HDAC6 Antibody [ARC0805] (A307941) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 10s.



Immunohistochemistry analysis of paraffin-embedded human breast cancer tissue using Anti-HDAC6 Antibody [ARC0805] (A307941) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunoprecipitation analysis of $300\mu g$ extracts of HeLa cells using $3\mu g$ of Anti-HDAC6 Antibody [ARC0805] (A307941). This Western blot was performed on the immunoprecipitate using Anti-HDAC6 Antibody [ARC0805] (A307941) at a dilution of 1:1000.