


Anti-IL-15 Antibody [ARC53689] (A308095)

Specifications:

Name:	Anti-IL-15 Antibody [ARC53689]
Description:	Rabbit monoclonal [ARC53689] antibody to IL-15.
Applications:	IHC
Recommended Dilutions:	IHC: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 49-162 of human IL15 (NP_000576.1).
Sequence:	NWVNVISDLKKIEDLIQSMHIDATLYTESDVHPSCKV TAMKCFLLLELQVISLES GDAS IHDTVENLILANNSLSSNGNVTESGCKECEEELEEKNIKEFLQSFVHIVQMFINTS
Host:	Rabbit
Clonality:	Monoclonal
Clone ID:	ARC53689
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.05% Proclin 300.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.


Images:




Immunohistochemistry analysis of paraffin-embedded human lung cancer using Anti-IL-15 Antibody [ARC53689] (A308095) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.

Anti-IL-15 Antibody [ARC53689] (A308095)


Images continued:



Immunohistochemistry analysis of paraffin-embedded human lung using Anti-IL-15 Antibody [ARC53689] (A308095) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded rat lung using Anti-IL-15 Antibody [ARC53689] (A308095) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse spleen using Anti-IL-15 Antibody [ARC53689] (A308095) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.