


Anti-PARP1 Antibody (A308167)

Specifications:

| | |
|------------------------|---|
| Name: | Anti-PARP1 Antibody |
| Description: | Rabbit polyclonal antibody to PARP1. |
| Applications: | WB, IHC, ICC/IF |
| Recommended Dilutions: | WB: 1:2,000-1:6,000, IHC: 1:50-1:200, ICC/IF: 1:50-1:200 |
| Reactivity: | Human, Mouse, Rat |
| Immunogen: | A synthetic peptide corresponding to a sequence within amino acids 200-300 of human PARP1 (NP_001609.2). |
| Sequence: | PGVKSEGKRKGDVVDGVDEVAKKKSKKEKDKDSKLEKALKAQNDLIWNIKDELKKVCS TNDLKELLIFNKQQVPSGESAILDRVADGMVFGALLPCEECSG |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Isotype: | IgG |
| Conjugate: | Unconjugated |
| Purification: | Affinity purification. |
| Molecular Weight: | 114 kDa |
| Product Form: | Liquid |
| Formulation: | Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal. |
| Storage: | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| Disclaimer: | This product is for research use only. It is not intended for diagnostic or therapeutic use. |


Images:




Western blot analysis of extracts from wild type (WT) and PARP1 knockout (KO) HeLa cells, using Anti-PARP1 Antibody (A308167) at 1:5,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 10s.

Anti-PARP1 Antibody (A308167)


Images continued:



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue using Anti-PARP1 Antibody (A308167) at a dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunofluorescence analysis of C6 cells using Anti-PARP1 Antibody (A308167) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of NIH-3T3 cells using Anti-PARP1 Antibody (A308167) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).