


## Anti-CD14 Antibody [ARC0353] (A308358)

### Specifications:


Name:	Anti-CD14 Antibody [ARC0353]
Description:	Rabbit monoclonal [ARC0353] antibody to CD14.
Applications:	WB, IHC, ICC/IF, Flow Cytometry
Recommended Dilutions:	WB: 1:500-1:2,000, IHC: 1:50-1:200, ICC/IF: 1:50-1:200, Flow Cytometry: 1:50-1:200
Reactivity:	Human, Mouse
Immunogen:	A synthetic peptide corresponding to a sequence within amino acids 250-350 of human CD14 (P08571).
Sequence:	QPHSLDLSHNSLRATVNPSPAPRCMWSSALNSLNLSFAGLEQVPKGLPAKLRVLDLSCN RLNRAPQPDELPEVDNLTLDGNPFLVPGTALPHEGSMNSGVVP
Host:	Rabbit
Clonality:	Monoclonal
Clone ID:	ARC0353
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	50 - 65 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

## Anti-CD14 Antibody [ARC0353] (A308358)


### Images:




Western blot analysis of extracts of various cell lines, using Anti-CD14 Antibody [ARC0353] (A308358) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 3min.



Immunohistochemistry analysis of paraffin-embedded human lung cancer using Anti-CD14 Antibody [ARC0353] (A308358) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunofluorescence analysis of THP-1 cells using Anti-CD14 Antibody [ARC0353] (A308358) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).



Flow cytometry analysis of A549 cells, stained with Rabbit IgG isotype control (10 µg/ml, blue line) or Anti-CD14 Antibody [ARC0353] (A308358), (10 µg/ml green line), followed by Alexa Fluor 647 conjugated goat anti-rabbit polyclonal antibody (1:600 dilution) staining. Non-fluorescently stained A549 cells, used as blank control (red line).