


Anti-hnRNP Q Antibody [ARC1656] (A308784)

Specifications:


Name:	Anti-hnRNP Q Antibody [ARC1656]
Description:	Rabbit monoclonal [ARC1656] antibody to hnRNP Q.
Applications:	WB, IHC, ICC/IF
Recommended Dilutions:	WB: 1:500-1:2,000, IHC: 1:50-1:200, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	A synthetic peptide corresponding to a sequence within amino acids 524-623 of human hnRNP Q/SYNCRIP (O60506).
Sequence:	SQRGGPGSARGVARGGAQQQRGRGVRGARGGRGGNVGGKRKADGYNQPDSKRRQTN NQNWGSQPIAQQPLQGGDHSGNYGYKSENQEFYQDTFGQQWK
Host:	Rabbit
Clonality:	Monoclonal
Clone ID:	ARC1656
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	70 - 85 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-hnRNP Q Antibody [ARC1656] (A308784)


Images:




Western blot analysis of extracts of various cell lines, using Anti-hnRNP Q Antibody [ARC1656] (A308784) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 3s.



Immunohistochemistry analysis of paraffin-embedded human liver cancer using Anti-hnRNP Q Antibody [ARC1656] (A308784) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse brain using Anti-hnRNP Q Antibody [ARC1656] (A308784) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunofluorescence analysis of C6 cells using Anti-hnRNP Q Antibody [ARC1656] (A308784) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).