


## Anti-CCR3 Antibody (A309001)

#### Specifications:

Name:	Anti-CCR3 Antibody
Description:	Rabbit polyclonal antibody to CCR3.
Applications:	WB, IHC
Recommended Dilutions:	WB: 1:500-1:1,000, IHC: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence within amino acids 146-226 of human CCR3 (NP_001828.1).
Sequence:	TVTFGVITSIVTWGLAVLAALPEFIFYETEELFEETLCSALYPEDTVYSWRHFHTLRM TIFCLVLPLLVMAICYTGIIKTL
Host:	Rabbit
Clonality:	Polyclonal
lsotype:	lgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	50 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

### Images:




Western blot analysis of extracts of various cell lines, using Anti-CCR3 Antibody (A309001) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 30s.


# antibodies

## Anti-CCR3 Antibody (A309001)

Images continued:



Immunohistochemistry analysis of paraffin-embedded rat spleen using Anti-CCR3 Antibody (A309001) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse spleen using Anti-CCR3 Antibody (A309001) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.