


## Anti-MyD88 Antibody [ARC52507 + ARC52509] (A309412)

### Specifications:

Name:	Anti-MyD88 Antibody [ARC52507 + ARC52509]
Description:	Rabbit Oligoclonal [ARC52507 + ARC52509] antibody to MyD88.
Applications:	WB
Recommended Dilutions:	WB: 1:2,000-1:9,000
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant protein of human MyD88.
Host:	Rabbit
Clonality:	Oligoclonal
Clone ID:	ARC52507 + ARC52509
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.05% Proclin 300.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.


### Images:



Western blot analysis of various lysates, using Anti-MyD88 Antibody [ARC52507 + ARC52509] (A309412) at 1:8,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 30s.

## Anti-MyD88 Antibody [ARC52507 + ARC52509] (A309412)

Images continued:



Western blot analysis of various lysates, using Anti-MyD88 Antibody [ARC52507 + ARC52509] (A309412) at 1:8,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 $\mu$ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Enhanced Kit (RM00021). Exposure time: 60s.