


Anti-TrkA +TRkB + TrkC Antibody [ARC2649] (A309687)

Specifications:

Name:	Anti-TrkA +TRkB + TrkC Antibody [ARC2649]
Description:	Rabbit monoclonal [ARC2649] antibody to TrkA +TRkB + TrkC.
Applications:	WB, IHC
Recommended Dilutions:	WB: 1:500-1:1,000, IHC: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	A synthesized peptide derived from human TrkA+B+C.
Host:	Rabbit
Clonality:	Monoclonal
Clone ID:	ARC2649
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	140 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.


Images:




Western blot analysis of extracts of various cell lines, using Anti-TrkA +TRkB + TrkC Antibody [ARC2649] (A309687) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 1s.

Anti-TrkA +TRkB + TrkC Antibody [ARC2649] (A309687)


Images continued:



Western blot analysis of extracts of SH-SY5Y cells, using Anti-TrkA +TRkB + TrkC Antibody [ARC2649] (A309687) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 180s.



Immunohistochemistry analysis of paraffin-embedded human brain tissue using Anti-TrkA +TRkB + TrkC Antibody [ARC2649] (A309687) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse brain using Anti-TrkA +TRkB + TrkC Antibody [ARC2649] (A309687) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.