


Anti-Claudin18 Antibody [ARC5062-02] (A309733)

Specifications:

| | |
|------------------------|---|
| Name: | Anti-Claudin18 Antibody [ARC5062-02] |
| Description: | Rabbit monoclonal [ARC5062-02] antibody to Claudin18. |
| Applications: | IHC |
| Recommended Dilutions: | IHC: 1:50-1:200 |
| Reactivity: | Human |
| Immunogen: | A synthetic peptide of human CLDN18.2. |
| Host: | Rabbit |
| Clonality: | Monoclonal |
| Clone ID: | ARC5062-02 |
| Isotype: | IgG |
| Conjugate: | Unconjugated |
| Purification: | Affinity purification. |
| Product Form: | Liquid |
| Formulation: | Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.05% Proclin 300. |
| Storage: | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| Disclaimer: | This product is for research use only. It is not intended for diagnostic or therapeutic use. |


Images:




Immunohistochemistry analysis of paraffin-embedded human esophageal n (negative control sample) using Anti-Claudin18 Antibody [ARC5062-02] (A309733) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.

Anti-Claudin18 Antibody [ARC5062-02] (A309733)


Images continued:



Immunohistochemistry analysis of paraffin-embedded human normal colon (negative control sample) using Anti-Claudin18 Antibody [ARC5062-02] (A309733) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human normal lung (negative control sample) using Anti-Claudin18 Antibody [ARC5062-02] (A309733) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human normal pancreas (negative control sample) using Anti-Claudin18 Antibody [ARC5062-02] (A309733) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.