


Anti-IL-1RA Antibody [ARC0524] (A80574)

Specifications:


Name:	Anti-IL-1RA Antibody [ARC0524]
Description:	Rabbit monoclonal [ARC0524] antibody to IL-1RA.
Applications:	WB, ICC/IF
Recommended Dilutions:	WB: 1:500-1:2,000, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 100-200 of human IL1RA (P18510).
Sequence:	ETRLQLEAVNITDLSNRKQDKRFAFIRSDSGPTTSFESAACPGWFLCTAMEADQPVS LTNMPDEGVMVTKFYFQEDE
Host:	Rabbit
Clonality:	Monoclonal
Clone ID:	ARC0524
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	22 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-IL-1RA Antibody [ARC0524] (A80574)


Images:




Western blot analysis of extracts of various cell lines, using Anti-IL-1RA Antibody [ARC0524] (A80574) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 1s.



Western blot analysis of extracts of various cell lines, using Anti-IL-1RA Antibody [ARC0524] (A80574) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 10s.



Confocal imaging of human skin using Anti-IL-1RA Antibody [ARC0524] (A80574), at a dilution of 1:100, (red). DAPI was used for nuclear staining (Blue). Objective: 60x.



Confocal imaging of mouse tail ear using Anti-IL-1RA Antibody [ARC0524] (A80574), at a dilution of 1:100, (red). DAPI was used for nuclear staining (Blue). Objective: 60x.