


Anti-PARK7 Antibody (A83672)

Specifications:

Name:	Anti-PARK7 Antibody
Description:	Goat polyclonal antibody to PARK7.
Applications:	ELISA, WB, IF
Reactivity:	Human, Mouse, Rat
Immunogen:	Synthetic peptide corresponding to Human PARK7 (C terminal).
Sequence:	C-AAQVKAPLVVKD
Host:	Goat
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Concentration:	100 µg at 0.5 mg/ml.
Product Form:	Liquid
Formulation:	Supplied in Tris Buffered Saline, pH 7.30, with 0.02% Sodium Azide and 0.5% BSA.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.


Images:




Anti-PARK7 Antibody (A83672) (0.001µg/ml) staining of Human Cerebellum (A), Human Frontal Cortex (B), Human Hippocampus (C) Mouse Fetal Brain (D), Mouse Brain (E) and Rat Brain (F) lysates (35µg protein in RIPA buffer). Detected by chemiluminescence.

Anti-PARK7 Antibody (A83672)


Images continued:



Anti-PARK7 Antibody (A83672) (0.001 μ g/ml) staining of HeLa (A) and Jurkat (B) lysates (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



Anti-PARK7 Antibody (A83672) (3.75 μ g/ml) staining of paraffin embedded Human Prostate. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



Anti-PARK7 Antibody (A83672) (3.75 μ g/ml) staining of paraffin embedded Human Thyroid. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.