

## Anti-ZNF131 Antibody (A87682)


### Specifications:

	Anti ZNIC121 Antihadu
Name:	Anti-ZNF131 Antibody
Description:	Rabbit polyclonal antibody to ZNF131.
Applications:	WB, ICC/IF
Recommended Dilutions:	WB: 1:200-1:2,000, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 80-240 of human ZNF131 (NP_003423.1).
Sequence:	MAFRHLIEFTYTAKLMIQGEEEANDVWKAAEFLQMLEAIKALEVRNKENSAPLEENTT GKNEAKKRKIAETSNVITESLPSAESEPVEIEVEIAEGTIEVEDEGIETLEEVASAKQ SVKYIQSTGSSDDSALALLADITSKYRQGDRKGQIKEDGCPSDPT
Host:	Rabbit
Clonality:	Polyclonal
lsotype:	lgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	80 - 85 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.
Storage:	Shipped at $4^{\circ}$ C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.
Molecular Weight: Product Form: Formulation: Storage:	80 - 85 kDa Liquid Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal. Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.


# antibodies

### Anti-ZNF131 Antibody (A87682)

#### Images:



Western blot analysis of extracts of various cell lines, using Anti-ZNF131 Antibody (A87682) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 5s.



Immunofluorescence analysis of U-2 OS cells using Anti-ZNF131 Antibody (A87682) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).