


Anti-VAPA Antibody (A89072)

Specifications:

| Name: | Anti-VAPA Antibody |
|------------------------|--|
| Description: | Rabbit polyclonal antibody to VAPA. |
| Applications: | WB, IHC, ICC/IF |
| Recommended Dilutions: | WB: 1:500-1:2,000, IHC: 1:50-1:200, ICC/IF: 1:50-1:100 |
| Reactivity: | Human, Mouse, Rat |
| Immunogen: | Recombinant fusion protein containing a sequence corresponding to amino acids 137-227 of human VAPA (NP_919415.2). |
| Sequence: | DKLNDMEPSKAVPLNASKQDGPMPKPHSVSLNDTETRKLMEECKRLQGEMMKLSEENR HLRDEGLRLRKVAHSDKPGSTSTASFRDNVTSP |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Isotype: | lgG |
| Conjugate: | Unconjugated |
| Purification: | Affinity purification. |
| Molecular Weight: | 28 kDa |
| Product Form: | Liquid |
| Formulation: | Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal. |
| Storage: | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| Disclaimer: | This product is for research use only. It is not intended for diagnostic or therapeutic use. |

Images:




Western blot analysis of extracts of various cell lines, using Anti-VAPA Antibody (A89072) at 1:3,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Enhanced Kit (RM00021). Exposure time: 90s.

antibodies

Anti-VAPA Antibody (A89072)

Images continued:



Immunohistochemistry analysis of paraffin-embedded human stomach using Anti-VAPA Antibody (A89072) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.