


Anti-TRPM2 Antibody (A9799)

Specifications:


Name:	Anti-TRPM2 Antibody
Description:	Rabbit polyclonal antibody to TRPM2.
Applications:	WB, ICC/IF
Recommended Dilutions:	WB: 1:100-1:500, ICC/IF: 1:50-1:100
Reactivity:	Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1204-1503 of human TRPM2 (NP_003298.1).
Sequence:	SSEADVPTLASQKAAEEPDAEPGGRKKTEEPGDSYHVNARHLLYPNCPVTRFRFPVNEK VPWETEFLLIYDPPFYTAERKDAAMDPMGDTLEPLSTIQYNVVDGLRDRRSFHGPTYV QAGLPLNPMGRTGLRGRGSLSCFGPNHTLYPMVTRWRRNEDGAICRKSIIKKMLEVLVV KLPLSEHWALPGGSREPGEMLPKLRILRQEHWPSFENLLKCGMEVYKGYMDDPRNT DNAWIETVAVSVHFQDQNDVELNRLNSNLHACDSGASIRWQVVDRIPLYANHKTLQ KAAAEFGAHY
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	171 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-TRPM2 Antibody (A9799)


Images:



Western blot analysis of extracts of Mouse lung, using Anti-TRPM2 Antibody (A9799) at 1:500 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Enhanced Kit (RM00021). Exposure time: 90s.



Immunofluorescence analysis of rat brain using Anti-TRPM2 Antibody (A9799) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of mouse brain using Anti-TRPM2 Antibody (A9799) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).