

## Anti-CD204 Antibody (A90674)


Specifications:

Name:	Anti-CD204 Antibody
Description:	Rabbit polyclonal antibody to CD204.
Applications:	WB
Recommended Dilutions:	WB: 1:500-1:1,000
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 192-451 of human MSR1 (NP_619729.1).
Sequence:	LNIENLNGKIQENTFKQQEEISKLEERVYNVSAEIMAMKEEQVHLEQEIKGEVKVLNN ITNDLRLKDWEHSQTLRNITLIQGPPGPPGEKGDRGPTGESGPRGFPGPIGPPGLKGD RGAIGFPGSRGLPGYAGRPGNSGPKGQKGEKGSGNTLTPFTKVRLVGGSGPHEGRVEI LHSGQWGTICDDRWEVRVGQVVCRSLGYPGVQAVHKAAHFGQGTGPIWLNEVFCFGRE SSIEECKIRQWGTRACSHSEDAGVTCTL
Host:	Rabbit
Clonality:	Polyclonal
Clonality: Isotype:	Polyclonal IgG
-	
Isotype:	lgG
Isotype: Conjugate:	IgG Unconjugated
Isotype: Conjugate: Purification:	IgG Unconjugated Affinity purification.
Isotype: Conjugate: Purification: Molecular Weight:	IgG Unconjugated Affinity purification. 60 - 90 kDa
Isotype: Conjugate: Purification: Molecular Weight: Product Form:	IgG Unconjugated Affinity purification. 60 - 90 kDa Liquid Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium

## antibodies

## Anti-CD204 Antibody (A90674)

Images:



Western blot analysis of extracts of various cell lines, using Anti-CD204 Antibody (A90674) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 90s.