


## Anti-ATP6V0C Antibody (A91538)

## Specifications:

Name:	Anti-ATP6V0C Antibody
Description:	Rabbit polyclonal antibody to ATP6V0C.
Applications:	WB, ICC/IF
Recommended Dilutions:	WB: 1:500-1:1,000, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human ATP6V0C (NP_001185498.1).
Sequence:	MSESKSGPEYASFFAVMGASAAMVFSALGAAYGTAKSGTGIAAMSVMRPEQIMKSIIP VVMAGIIAIYGLVVAVLIANSLNDDISLYKSFLQLGAGLSVG
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	lgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	16 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.05% Proclin 300.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.
Discialmer.	This product is for research use only. It is not intended for diagnostic of therapeutic use.

## Images:




Western blot analysis of various lysates, using Anti-ATP6V0C Antibody (A91538) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 60s.



## Anti-ATP6V0C Antibody (A91538)

Images continued:



Immunofluorescence analysis of MCF7 cells using Anti-ATP6V0C Antibody (A91538) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).