


## Anti-PEDF Antibody (A93125)

### Specifications:

|                        |  |
|------------------------|--|
| Name:                  | Anti-PEDF Antibody   |
| Description:           | Rabbit polyclonal antibody to PEDF.  |
| Applications:          | WB, ICC/IF   |
| Recommended Dilutions: | WB: 1:500-1:2,000, ICC/IF: 1:50-1:200  |
| Reactivity:            | Human, Mouse, Rat  |
| Immunogen:             | A synthetic peptide corresponding to a sequence within amino acids 150-250 of human PEDF/SERPINF1 (NP_002606.3). |
| Sequence:              | IKSSFVAPLEKSYGTRPRVLTGNPRLDLQEINNWWQAQMKGKRLARSTKEIPDEISILL<br>LGVAHFQGWVTKFDSRKSLEDFYLDEERTVRVPMMSDPKAV         |
| Host:                  | Rabbit   |
| Clonality:             | Polyclonal   |
| Isotype:               | IgG  |
| Conjugate:             | Unconjugated   |
| Purification:          | Affinity purification.   |
| Molecular Weight:      | 46 kDa   |
| Product Form:          | Liquid   |
| Formulation:           | Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.                           |
| Storage:               | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.                            |
| Disclaimer:            | This product is for research use only. It is not intended for diagnostic or therapeutic use.                     |

### Images:



Western blot analysis of extracts of various cell lines, using Anti-PEDF Antibody (A93125) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 30s.