

Anti-Myeloid leukemia factor 1 Antibody (A10262)

Specifications:

Name: Anti-Myeloid leukemia factor 1 Antibody

Description: Rabbit polyclonal antibody to Myeloid leukemia factor 1.

Applications: WB, ICC/IF

Recommended Dilutions: WB: 1:500-1:2,000, ICC/IF: 1:50-1:100

Reactivity: Human, Mouse

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-268 of

human MLF1 (NP_071888.1).

Sequence: MFRMLNSSFEDDPFFSESILAHRENMRQMIRSFSEPFGRDLLSISDGRGRAHNRRGHN

DGEDSLTHTDVSSFQTMDQMVSNMRNYMQKLERNFGQLSVDPNGHSFCSSSVMTYSKI GDEPPKVFQASTQTRRAPGGIKETRKAMRDSDSGLEKMAIGHHIHDRAHVIKKSKNKK TGDEEVNQEFINMNESDAHAFDEEWQSEVLKYKPGRHNLGNTRMRSVGHENPGSRELK

RREKPQQSPAIEHGRRSNVLGDKLHIKGSSVKSNKK

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Conjugate: Unconjugated

Purification: Affinity purification.

Molecular Weight: 35 kDa

Product Form: Liquid

Formulation: Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium

Azide.


Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.




Anti-Myeloid leukemia factor 1 Antibody (A10262)

Images:



Western blot analysis of extracts of various cell lines, using Anti-Myeloid leukemia factor 1 Antibody (A10262) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 90s.



Immunofluorescence analysis of A549 cells using Anti-Myeloid leukemia factor 1 Antibody (A10262).