


Anti-STAT6 (phospho Tyr641) Antibody (A10952)

Specifications:

Name:	Anti-STAT6 (phospho Tyr641) Antibody
Description:	Rabbit polyclonal antibody to STAT6 (phospho Tyr641).
Applications:	WB, IHC
Recommended Dilutions:	WB: 1:500-1:2,000, IHC: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	A synthetic phosphorylated peptide around Y641 of human STAT6 (NP_003144.3).
Sequence:	RGYVP
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	110 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.05% Proclin 300.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.


Images:




Western blot analysis of extracts from HeLa cells, using Anti-STAT6 (phospho Tyr641) Antibody (A10952). The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% BSA.

Anti-STAT6 (phospho Tyr641) Antibody (A10952)


Images continued:



Immunohistochemistry analysis of paraffin-embedded human stomach using Anti-STAT6 (phospho Tyr641) Antibody (A10952) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse kidney using Anti-STAT6 (phospho Tyr641) Antibody (A10952) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse spleen using Anti-STAT6 (phospho Tyr641) Antibody (A10952) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.