

Anti-Dynamin 2 Antibody (A12643)

Specifications:

Name: Anti-Dynamin 2 Antibody

Description: Rabbit polyclonal antibody to Dynamin 2.

Applications: WB, ICC/IF

Recommended Dilutions: WB: 1:500-1:2,000, ICC/IF: 1:50-1:200

Reactivity: Human, Mouse, Rat

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 607-866

of human DNM2 (NP_004936.2).

Sequence: EDVDSWKASFLRAGVYPEKDQAENEDGAQENTFSMDPQLERQVETIRNLVDSYVAIIN

KSIRDLMPKTIMHLMINNTKAFIHHELLAYLYSSADQSSLMEESADQAQRRDDMLRMY HALKEALNIIGDISTSTVSTPVPPPVDDTWLQSASSHSPTPQRRPVSSIHPPGRPPAV RGPTPGPPLIPVPVGAAASFSAPPIPSRPGPQSVFANSDLFPAPPQIPSRPVRIPPGI

PPGVPSRRPPAAPSRPTIIRPAEPSLLD

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Conjugate: Unconjugated

Purification: Affinity purification.

Molecular Weight: 110 kDa

Product Form: Liquid

Formulation: Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium

Azide.


Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.




Anti-Dynamin 2 Antibody (A12643)


Images:




Western blot analysis of extracts of various cell lines, using Anti-Dynamin 2 Antibody (A12643) at 1:500 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 90s.



Immunofluorescence analysis of U2OS cells using Anti-Dynamin 2 Antibody (A12643) at a dilution of 1:150. DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of NIH/3T3 cells using Anti-Dynamin 2 Antibody (A12643) at a dilution of 1:150. DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of PC-12 cells using Anti-Dynamin 2 Antibody (A12643) at a dilution of 1:150. DAPI was used to stain the cell nuclei (blue).