

Anti-KPNA1 Antibody (A13627)


Specifications:

| mino acids 1-300 of |
|--|
| (RRNVATAEEETE (FRKLLSKEPNP /IQAGAVPIF (QNRLTMTRNA SYLSDGPNDKIQ |
| |
| |
| |
| |
| |
| |
| |
| 0.02% Sodium |
| / thaw cycles. |
| therapeutic use. |
| |


antibodies

Anti-KPNA1 Antibody (A13627)


Images:



Western blot analysis of extracts of various cell lines, using Anti-KPNA1 Antibody (A13627) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST.



Immunohistochemistry analysis of paraffin-embedded human lung cancer using Anti-KPNA1 Antibody (A13627) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human esophagus using Anti-KPNA1 Antibody (A13627) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.