


Anti-TNFAIP3 Antibody (A13892)

Specifications:


Name:	Anti-TNFAIP3 Antibody
Description:	Rabbit polyclonal antibody to TNFAIP3.
Applications:	WB, IHC
Recommended Dilutions:	WB: 1:100-1:500, IHC: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 600-738 of human TNFAIP3 (NP_006281.1).
Sequence:	DRTGTSKCRKAGCVYFGTPENKGFCTLCFIEYRENKHFAAASGKVSPTASRFQNTIPC LGRECGTLGSTMFEQYCYQKCFIEAQNQRFHEAKRTEEQLRSSQRRDVPRTTQSTSRPK CARASCKNILACRSEELCMECQH
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	90 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.05% Proclin 300.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-TNFAIP3 Antibody (A13892)


Images:




Western blot analysis of extracts of various cell lines, using Anti-TNFAIP3 Antibody (A13892) at 1:500 dilution. THP-1 cells were treated by PMA/TPA (200 nM) at 37°C for 15 minutes after serum-starvation overnight. THP-1 cells were treated by LPS (1 µg/ml) at 37°C for 8 hours. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 40s.



Immunohistochemistry analysis of paraffin-embedded human colon tissue using Anti-TNFAIP3 Antibody (A13892) at a dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human lung cancer using Anti-TNFAIP3 Antibody (A13892) at a dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded rat lung using Anti-TNFAIP3 Antibody (A13892) at a dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.